

### I-MASK+

# PROPHYLAXIS & EARLY OUTPATIENT TREATMENT PROTOCOL FOR COVID-19

Page 1/2

#### PROPHYLAXIS PROTOCOL

Ivermectin Prophylaxis for high risk individuals

0.2 mg/kg\* – one dose on day 1 and day 3, then take

one dose every 4 weeks

Post COVID-19 exposure prophylaxis\*\*
0.2 mg/kg\* – one dose on day 1 and day 3

Vitamin D3 1,000-3,000 IU/day

Vitamin C 1,000 mg twice a day

Quercetin 250 mg/day Zinc 50 mg/day

Melatonin 6 mg before bedtime (causes drowsiness)

#### **EARLY OUTPATIENT PROTOCOL\*\*\***

Ivermectin 0.2 mg/kg\* – one dose on day 1 and day 3

Vitamin D3 4,000 IU/day

Vitamin C 2,000 mg 2-3 times daily

Quercetin 250 mg twice a day

Zinc 100 mg/day

Melatonin 10 mg before bedtime

Aspirin 325 mg/day (unless contraindicated)

- \* ≈ 0.09 mg/lb per dose (take on an empty stomach with water). Example for a person of 60 kg (body weight): 60 kg × 0.2 mg = 12 mg please see conversion table (kg→lbs) on page 2 to calculate the appropriate ivermectin dose.
- \*\* To use if a household member is COVID-19 positive, or you have prolonged exposure to a COVID-19 positive patient without wearing a mask
- \*\*\* For late phase <u>hospitalized</u> patients see the FLCCC's **MATH+ Hospital Treatment Protocol for COVID-19** on www.flccc.net

Please regard our disclaimer and further information on page 2 of this document.





#### **WEAR MASKS**

Must wear cloth, surgical, or N95 mask (without valve) in all indoor spaces with nonhousehold persons.

Must wear a N95 mask (without valve) during prolonged exposure to nonhousehold persons in any confined, poorly ventilated area.



#### **KEEP DISTANCE**

Until the end of the Covid-19 crisis, we recommend keeping a minimum distance of approx. 2 m / 6 feet in public from people who are not from your own household.



#### **WASH HANDS**

We recommend, after a stay during and after outings from home (shopping, subway etc.), a thorough hand cleaning (20–30 sec. with soap), or also to use a hand disinfectant in between.



### I-MASK+

# PROPHYLAXIS & EARLY OUTPATIENT TREATMENT PROTOCOL FOR COVID-19

Page 2/2

#### **IVERMECTIN**

## Body weight conversion (kg/lb) for ivermectin dose in prophylaxis and treatment of COVID-19

Dose		Body weight	
$0.2 \text{ mg/kg} \approx 0.09 \text{ mg/lb}$ (Each tablet = 3 mg; doses rounded		Conversion (1kg ≈ 2.2 lbs) (doses calculated per	
(3 tablets=9 mg)	8 mg	32-40 kg	70-90 lb
(3.5 tablets)	10 mg	41-50 kg	91-110 lb
(4 tablets)	12 mg	51-59 kg	111-130 lb
(4.5 tablets)	13.5 mg	60-68 kg	131-150 lb
(5 tablets)	15 mg	69-77 kg	151-170 lb
(5.5 tablets)	16 mg	78-86 kg	171-190 lb
(6 tablets)	18 mg	87-95 kg	191-210 lb
(7 tablets = 21 mg)	20 mg	96-104 kg	211-230 lb
(7.5 tablets=22.5 mg)	22 mg	105-113 kg	231-250 lb
(8 tablets)	24 mg	114-122 kg	251-270 lb
(9 tablets=27 mg)	26 mg	123-131 kg	271-290 lb
(9.5 tablets=28.5 mg)	28 mg	132-140 kg	291-310 lb

#### **DISCLAIMER**

The I-Mask+ Prophylaxis & Early Outpatient Treatment Protocol for COVID-19 and the MATH+ Hospital Treatment Protocol for COVID-19 are solely for educational purposes regarding potentially beneficial therapies for COVID-19. Never disregard professional medical advice because of something you have read on our website and releases. It is not intended to be a substitute for professional medical advice, diagnosis, or treatment in regards to any patient. Treatment for an individual patient should rely on the judgement of your physician or other qualified health provider. Always seek their advice with any questions you may have regarding your health or medical condition.

A summary of the published data supporting the rationale for Ivermectin use in our I-MASK+ protocol can be downloaded from www.flccc.net/i-mask-prophylaxis-treatment-protocol/

For updates, references, and information on the FLCCC Alliance, the I-Mask+ Prophylaxis & Early Outpatient Treatment Protocol for COVID-19 and the MATH+ Hospital Treatment Protocol for COVID-19, please visit our website.